

GENERALIZED STATISTICS AND DYNAMICS IN CURVED SPACETIME

V.Bardek, S.Meljanac and A.Perica

Rudjer Bošković Institute, P.O.B. 1016,

41001 Zagreb, Croatia

(January 4, 1994)

Abstract

We consider the generalized momentum-dependent quon algebra in a dynamically evolving curved spacetime and perform a type of analysis similar to that of J.W.Goodison and D.J.Toms. We find that, at least in principle, all kinds of statistics may occur in some regions, i.e. phases in momentum space, depending on Bogoliubov coefficients determined by a specific dynamical model.

PACS numbers: 05.30. - d , 04.60. + n

The problem of allowed statistics, such as Bose, Fermi, para-Bose, para-Fermi, anyonic, infinite-quonic, and their role in physics is of great importance [1-4]. It has recently been claimed [5] that, starting with the global quon algebra [2] no generalized statistics is compatible with the dynamics in curved spacetime. The only allowed possibilities found are the ordinary Bose and Fermi statistics. However, the initial assumption that the parameter of deformation q is a real constant is too restrictive. Namely, two of us [6] proposed a more general quon algebra interpolating between all types of statistics simultaneously, and having the same basic properties as the global quon algebra [2].

In this Letter, starting with the generalized quon algebra, we classify and discuss possible statistics compatible with gravitationally induced particle creation in a dynamically evolving universe.

The generalized quon algebra [6] is given by

$$a_i a_j^\dagger - q_{ij} a_j^\dagger a_i = \delta_{ij}, \forall i, j \in S \quad (1)$$

Here $q_{ij} = q_{ji}^*$ are complex functions and S is a set of (discrete or continuous) 3-dimensional momenta. Norms in the Fock space are positive definite if $|q_{ij}| \leq 1$, for all pairs $i, j \in S$. If $|q_{ij}| < 1$, there are no commutation relations between a_i, a_j (or $a_i^\dagger, a_j^\dagger$) operators. However, in the exact limit when $|q_{ij}| = 1$ for every pair of indices $i, j \in S$, the following commutation relations are valid:

$$a_i a_j - q_{ij}^* a_j a_i = 0, \forall i, j \in S \quad (2)$$

In this case the number operator is $N_i = a_i^\dagger a_i$.

We consider generalized quon statistics in a dynamically evolving curved space-time which, prior to some initial time t_1 and subsequent to some later time t_2 , is flat. The spacetime is dynamic for $t_1 \leq t \leq t_2$ and given by the spatially flat Robertson-Walker metric [7]. In the first flat region where $t < t_1$ (in-region) we assume that annihilation and creation operators $a_i, a_i^\dagger$ satisfy the generalized quon algebra, Eq.(1). In the second flat region where $t > t_2$ (out-region) we assume that the corresponding annihilation and creation operators $b_i, b_i^\dagger$ satisfy the generalized quon algebra:

$$b_i b_j^\dagger - q'_{ij} b_j^\dagger b_i = \delta_{ij}, \forall i, j \in S. \quad (3)$$

We have denoted these operators by b_i and $b_i^\dagger$ because they, in general, differ from the corresponding one in the in-region. The general relation between the two sets of in- and out- operators is given by

$$a_i = \sum_j (\alpha_{ji} b_j + \beta_{ji}^* b_j^\dagger), \quad (4)$$

where α_{ji}, β_{ji} are Bogoliubov coefficients for the expanding universe [5],[7]. However, for the spacetime with the Robertson-Walker metric between the in- and out-regions, $t_1 < t < t_2$, the Bogoliubov coefficients are diagonal [7]. Therefore, we restrict ourselves to the case where

$$\begin{aligned}\alpha_{ij} &= \alpha_i \delta_{ij}, \\ \beta_{ij} &= \beta_i \delta_{ij}.\end{aligned}\tag{5}$$

Substituting (4,5) and its complex conjugate into (1), we easily obtain

$$\begin{aligned}\alpha_i \alpha_j^* (b_i b_j^\dagger - q_{ij} b_j^\dagger b_i) + \beta_i^* \alpha_j^* (b_i^\dagger b_j^\dagger - q_{ij} b_j^\dagger b_i^\dagger) + \alpha_i \beta_j (b_i b_j - q_{ij} b_j b_i) \\ + \beta_i^* \beta_j (b_i^\dagger b_j - q_{ij} b_j b_i^\dagger) = \delta_{ij}.\end{aligned}\tag{6}$$

Taking the expectation value of (6) with a vacuum state in the out-region yields,

$$|\alpha_i|^2 - q_{ii} |\beta_i|^2 = 1, \forall i \in S.\tag{7}$$

From (7) it follows that (6) can be reduced to

$$\begin{aligned}\alpha_i \beta_j (b_i b_j - q_{ij} b_j b_i) + \beta_i^* \alpha_j^* (b_i^\dagger b_j^\dagger - q_{ij} b_j^\dagger b_i^\dagger) + (1 - q_{ij} q'_{ji}) \beta_i^* \beta_j b_i^\dagger b_j + \\ (q'_{ij} - q_{ij}) \alpha_i \alpha_j^* b_j^\dagger b_i = 0.\end{aligned}\tag{8}$$

It is easy to see that the remaining relations following from (8) are

$$\begin{aligned}\alpha_i \beta_j (b_i b_j - q_{ij} b_j b_i) &= 0, \forall i, j \in S \\ \beta_i^* \alpha_j^* (b_i^\dagger b_j^\dagger - q_{ij} b_j^\dagger b_i^\dagger) &= 0, \forall i, j \in S \\ (1 - q_{ij} q'_{ji}) \beta_i^* \beta_j &= 0, i \neq j \\ (q'_{ij} - q_{ij}) \alpha_i \alpha_j^* &= 0, i \neq j \\ (q_{ii} - q'_{ii}) |\alpha_i|^2 &= (1 - q_{ii} q'_{ii}) |\beta_i|^2, \forall i \in S.\end{aligned}\tag{9}$$

Now, according to the possible values of Bogoliubov coefficients, we shall classify and discuss all possible solutions of the conditions (7), (9).

Solution (i): $\alpha_i \beta_i \neq 0, \forall i \in S$

From the first of Eqs. (9) it follows that

$$b_i b_j - q_{ij} b_j b_i = 0, \forall i, j \in S \quad (10)$$

and its hermitian conjugate. These commutation relations between b_i, b_j can exist only if $|q_{ij}| = 1$ for $\forall i, j \in S$, [6].

From Eq. (2) it follows that $q_{ij}^* = q_{ij}$. Hence, $q_{ij}^2 = 1$, i.e., $q_{ij} = \pm 1, \forall i, j \in S$. Furthermore, from the fourth and fifth of Eqs. (9) one obtains $q'_{ij} = q_{ij}$. Let us summarize solution (i): $q'_{ij} = q_{ij}, q_{ij}^2 = 1$ for every pair i,j.

Let us further discuss two possible cases. First, if the index i is a continuous variable, then the functions α_i, β_i are continuous functions of i. It follows from Eq. (7) that $q_{ii} = \text{const.}$, i.e., $q_{ii} = +1, \forall i \in S$ or $q_{ii} = -1, \forall i \in S$. There are no additional restrictions on q_{ij} if $i \neq j$. Note that, in the above considerations, q_{ij} has not necessarily been a continuous function. (However, if we assume that q_{ij} is also a continuous function in the variables i,j then $q_{ij} = \text{const.} = +1 \text{ or } -1, i, j \in S$.) Note that q_{ii} would have jumps if β_i could vanish.

The second possibility assumes that in addition to the continuous variable i there is a discrete variable $\mu = 1, \dots, p$. All the conditions (Eqs.(7), (9)) remain the same under the following substitutions: $\alpha_i, \beta_i \rightarrow \alpha_i^\mu, \beta_i^\mu, a_i, b_i \rightarrow a_i^\mu, b_i^\mu$, and $q_{ij} \rightarrow q_{i\mu, j\nu}$. If we assume that $q_{i\mu, j\nu}$ is a continuous function in the i,j variables, then $q_{i\mu, j\nu}$ reduces to $q_{\mu\nu}$. Solution (i) reads: $q_{\mu\mu} \neq \text{const.}, q'_{\mu\nu} = q_{\mu\nu}, q_{\mu\nu}^2 = 1, \mu, \nu = 1, \dots, p$.

Let us write the corresponding algebra:

$$\begin{aligned} a_i^\mu a_j^{\nu\dagger} - q_{\mu\nu} a_j^{\nu\dagger} a_i^\mu &= \delta_{ij} \delta_{\mu\nu}, \forall \mu, \nu = 1, \dots, p \\ a_i^\mu a_j^\nu - q_{\mu\nu} a_j^\nu a_i^\mu &= 0, \forall i, j \in S. \end{aligned} \quad (11)$$

This is a generalization of Green's oscillator algebra [1]. In the special case $q_{\alpha\beta} = \pm(2\delta_{\alpha\beta} - 1)$, one obtains p Green's oscillators of the Bose (Fermi) type for the upper (lower) sign. Hence, para-Bose and para-Fermi statistics are also allowed, in addition to the ordinary Bose and Fermi statistics.

Solution (ii): $\alpha_i \neq 0, \beta_i = 0, \forall i \in S$

Equation (7) gives $|\alpha_i| = 1, i \in S$ and the fourth and the fifth of Eqs. (9) give $q'_{ij} = q_{ij}$. In this case there are no restrictions on statistics. We can have arbitrary q_{ij} , with $|q_{ij}| \leq 1$ and $q'_{ij} = q_{ij}$. In this case anyons in a multivalued picture [8] are also allowed.

Solution (iii): $\alpha_i = 0, \beta_i \neq 0, \forall i \in S$

From Eq. (7) it follows that $|\beta_i|^2 = 1$ and $q_{ii} < 0, \forall i \in S$. Since $\beta_i \beta_j^* \neq 0$, the third of Eqs. (9) gives $q'^*_{ij} = q_{ij}^{-1}$. In addition, $|q'_{ij}| \leq 1$ and $|q_{ij}| \leq 1$, therefore $|q'_{ij}| = |q_{ij}| = 1$. Hence, we find that $q'_{ij} = q_{ij} = e^{i\Delta_{ij}}, i \neq j$, where $\Delta_{ij} = -\Delta_{ji}$ is a real function. For $i = j$, $q'_{ii} = q_{ii}^{-1}$, and in the same way as above we find that $q'_{ii} = q_{ii} = \pm 1$. Since $q_{ii} < 0$, the only possibility is $q'_{ii} = q_{ii} = -1$, which means the hard-core condition for all oscillators: $a_i^2 = b_i^2 = 0, \forall i \in S$.

Let us summarize solution (iii):

$q'_{ij} = q_{ij} = e^{i\Delta_{ij}}, i \neq j$ and $q'_{ii} = q_{ii} = -1, \forall i \in S$. It corresponds to anyons with the hard-core condition in a single-valued picture [9].

General solution (iv):

Solutions (i), (ii) and (iii) are characterized by the respective restriction s: $\alpha_i \beta_i \neq 0, \forall i \in S$; $\alpha_i \neq 0, \beta_i = 0, \forall i \in S$; $\alpha_i = 0, \beta_i \neq 0, \forall i \in S$. According to [5],[7], solution (ii) is not physically acceptable. In a similar way, solution (iii) seems to be not allowed unless there is a special dynamical reason for that. So we are left with solution (i) allowing the Bose, Fermi, and generalized Green's oscillator algebra to be consistent with the dynamical evolution of spacetime.

However, even solution (i), might not be realistic. Therefore, we now assume no restrictions on α_i, β_i , except the necessary condition Eq.(7). Let us decompose the initial set of indices i into three not overlapping subsets:

$$\begin{aligned} M_1 &= \{i \in S, \alpha_i \beta_i \neq 0\}, \\ M_2 &= \{i \in S, \beta_i = 0\}, \\ M_3 &= \{i \in S, \alpha_i = 0\}. \end{aligned} \tag{12}$$

Note that $M_1 \cup M_2 \cup M_3 = S$.

It is obvious that solution (1) applies to set M_1 , solution (ii) to set M_2 , and solution (iii) to set M_3 . Equations (9) with indices i, j from different subsets (M_1, M_2) , (M_1, M_3) , (M_2, M_3) , are satisfied by $q'_{ij} = q_{ij} = \pm 1$. Hence the general solution for α_i, β_i , calculated in a dynamical model of evolving spacetime, allows all types of statistics, at least in principle. The main conclusion is that $q'_{ij} = q_{ij}$, i.e., that q_{ij} cannot be changed in the course of dynamical evolution. The question of existence, structure, and topology of subsets M_2, M_3 is deeply connected with a concrete dynamical model of evolving spacetime, and is beyond the scope of this paper.

Nevertheless, let us assume that nontrivial sets M_2, M_3 exist. Hence, there may

exist several different regions (i.e., "phases" in momentum space) with solutions M_1 of the Bose or Fermi type. The regions are separated by a boundary of the type M_2 or M_3 . All types of statistics (including infinite-quonic statistics) are possible on the $M_2^{\beta=0}$ boundary. Only anyons with the hard-core condition are possible on the $M_3^{\beta=0}$ boundary. Transition from one M_1 phase to another M_1 phase through $M_3^{\beta=0}$ may change the Bose-type to Fermi-type of statistics. However, transition through $M_3^{\alpha=0}$ is possible only if the neighboring phases are of the Fermi-type and, hence, cannot change the type of statistics.

Finally, let us mention that our analysis, although general, is still restricted to the generalized quon algebra (Eq.(1)). However, there are other deformed algebras [10], not included in Eq.(1), for which a similar analysis can be performed.

This work was supported by the Scientific Fund of the Republic of Croatia.

References

- [1] H.S.Green, Phys.Rev.**90**, 270 (1953).
Y.Ohnuki and S.Kamefuchi, *Quantum Field Theory and Parastatistics* (University of Tokyo Press, Tokyo, Springer, Berlin, 1982).
- [2] O.W.Greenberg, Phys.Rev.Lett.**64**, 705 (1990), Phys.Rev.D**43**, 4111 (1991).
- [3] O.W.Greenberg, D.M.Greenberger, and T.V.Greenberger, in *Proceedings of the International Conference on Fundamental Aspects of Quantum Theory*, 1993.
A.Strominger, Phys.Rev.Lett.**71**, 3397 (1993).
- [4] J.M.Leinaas and J.Myrheim, Nuovo Cim.**37B**, 1 (1977). F.Wilczek, Phys.Rev.Lett.**48**, 114 (1982).
- [5] J.W.Goodison and D.J.Toms, Phys.Rev.Lett.**71**, 3240 (1993).
- [6] S.Meljanac and A.Perica, *Bose, Fermi, para-Bose, para-Fermi, and anyon statistics: The R-matrix approach to deformed associative algebras.*, submitted to Phys.Rev.Lett.
- [7] L.Parker, Phys.Rev.**183**, 1057 (1969), Phys.Rev.D**3**, 346 (1971).
N.D.Birrell and P.C.W.Davies, *Quantum fields in curved space* (Cambridge University Press, 1982).
- [8] V.Bardek, M.Doresic, and S.Meljanac, *Anyons as quon particles*, to be published in Phys.Rev.D.

- [9] A.Lerda and S.Sciuto, Nucl.Phys.B**401**, 613 (1993).
- [10] W.Pusz and S.Woronowicz, Rep.Math.Phys.**27**, 231 (1989). D.B.Fairlie and C.K.Zachos, Phys.Lett.B**256**, 43 (1991).